

# Biomechanical Considerations and Occlusal Schemes in Implant Restorations; A Review, Part I: Fixed Protheses

Safoura Ghodsi <sup>1</sup>, Arezoo Mazaheri Nazarifar <sup>2</sup>, Sasan Rasaeipour <sup>3</sup>, Azadeh Sadeghi <sup>4</sup> 

<sup>1</sup> Associate professor, Dental Research Center, Dentistry Research Institute, Department of Prosthodontics, School of Dentistry, Tehran University of Medical Sciences, Tehran, Iran

<sup>2</sup> Postgraduate Student, Department of Prosthodontics, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>3</sup> Assistant Professor, Department of Prosthodontics, School of Dentistry, Tehran University of Medical Sciences, Tehran, Iran

<sup>4</sup> Postgraduate Student, Department of Prosthodontics, School of Dentistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran

## Abstract

**Background and Aim:** The aim of this study was to gather available scientific data on appropriate occlusal scheme in each type of fixed implant restoration with the least adverse effect on implant and peri-implant tissues.

**Materials and Methods:** An extensive search was performed in PubMed, Scopus, Embase, and Google Scholar using related keywords. Studies related to evaluation of different occlusal schemes, and occlusion in implant dentistry were selected, reviewed, and discussed.

**Results:** The selected keywords yielded 995 search results in PubMed, 417 in Embase, and 500 in Scopus. After duplicate removal and title/abstract analysis, 83 studies were selected for full-text review. Finally, 43 studies met the requirement of inclusion/exclusion criteria and were included.

**Conclusion:** Selecting the right occlusal scheme for implant restorations is complex. Clinicians must consider multiple factors to prevent occlusal overloading. Following scientific guidelines ensures long-term success and predictability in implant-supported protheses.

**Key Words:** biomechanics, implant occlusion, occlusal considerations, occlusal scheme, prosthetic considerations

 Corresponding author:  
Azadeh Sadeghi, Postgraduate Student, Department of Prosthodontics, School of Dentistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran

azadesadeghi137@yahoo.com

Received: 2 April 2023

Accepted: 28 June 2023

➤ **Cite this article as:** Ghodsi S, Mazaheri Nazaheri A, Rasaeipour S, Sadeghi A. Biomechanical Considerations and Occlusal Schemes in Implant Restorations; A Review, Part I: Fixed Protheses. *J Iran Dent Assoc.* 2023; 35(1-2):21-31. doi: 10.34172/jida.A-10-1-991

## Introduction

Implant-based treatments are highly desirable for replacing missing teeth due to their high predictability and success rates. These treatments boast a success rate of over 80% for up to 16 years (1-4). Occlusion is an important factor in successful long-term serviceability of dental protheses (1-3,5,6). Considering the lack of periodontal cushioning effect in implants, occlusion could even play a more important role

in implant-based treatments, and in fact, is one of the most important determining factors in long run implant success (1,6) The effect of occlusion on long-term success returns to implant sensitivity to biomechanical stresses. (1-6) Occlusal stress has been suggested as one of the primary causes of biomechanical (e.g. screw loosening, implant component fracture, detachment of implant crowns ) and biological (e.g. marginal bone loss) implant complications

(1,5-9). The role of occlusal overloading in biological implant failures is controversial. While animal studies have not confirmed this relationship, clinical studies have shown a clear correlation between parafunction and implant marginal bone loss (10-13). Applying correct occlusal patterns can assist clinicians in delivering long lasting and predictable treatments for their patients (1-3,5-10). Well recognized occlusal schemes with some modifications can be applied in implant dentistry (1-3,5-10) These modifications address implant sensitivity to lateral forces, lack of proprioception as a warning alarm in traumatic occlusal contacts in implants, and mechanical properties of dental implant systems (3,14,5-10) Since there is no acceptable codified conclusion available on proper occlusion types in fixed implant restorations, the present review study aims to suggest evidence-based occlusal schemes for different types of implant-supported fixed prostheses based on diverse scientific documents available.

### Materials and Methods

A comprehensive electronic search was made using PubMed, Medline, Scopus, Embase, and Google Scholar from 1887 to 2020. The following keywords, with different combinations were searched in article titles, abstracts, or keywords: occlusal consideration, occlusal scheme, occlusal pattern, occlusion, occlus\*, guidance, biomechanic, mechanical factor, dental implant, implant, and fixed prosthes\*. Using reference management software (Endnote X8; Thomson Reuters), duplicated studies were eliminated, and articles were selected based on title-abstract analysis, followed by full text evaluation. Two independent reviewers evaluated the studies. In cases of disagreement, they engaged in discussions until consensus was reached. The included studies comprised peer-reviewed articles in the form of review articles, cohort studies, case-control studies, cross-sectional studies, case series or reports, and experimental studies. The included articles were selected in a way that they discussed occlusion in fixed implant prostheses, the consequences of

occlusion and biomechanical forces on dental implants, and studies on stress profiles or occlusion principles in implant restorations. The studies focusing on removable restorations or tooth-supported prostheses were excluded. Animal studies were not also taken into consideration. The Data on the recommended occlusal schemes, and biomechanical principles and guidelines were extracted.


### Results

The search results for the selected keywords yielded 995 published articles in PubMed, 417 in Embase, and 500 in Scopus. After duplicate removal and title/abstract analysis, 83 studies were selected for full-text review. Finally, 43 studies met the requirement of inclusion/exclusion criteria and were included. Detailed information on search process is summarized in figure 1.

The articles showed extensive diversity in preferred occlusion for certain implant prostheses. To address this, the present study reviewed various suggested occlusal schemes and proposed the most prevalent or acceptable one. For proposing the most acceptable occlusion in such cases, the authors discussed and chose rational scheme based on the implant occlusion principles, and available finite element analysis or experimental studies which focused on stress distribution profile. Considering limited number of studies on this basic scientific field, the authors tried to subsume all the available evidence for implant-fixed prosthesis occlusion; although some of the evidences were not strong enough to be conclusive. The lack of adequate or strong scientific evidences for supporting specific occlusal scheme in some prosthetic type has been notified as "limited information". These areas call for further research.

### Discussion

Several occlusal schemes have been introduced for natural dentition, each with its own advantages and indications. These schemes will be summarized here based on the latest published terminology, as understanding this preliminary information aids in comprehending



**Figure 1.** The study selection and identification chart

the content in a coordinated manner. Considering the specific requirements of implants, some of these patterns can be adapted for dental implants with certain modifications (3,14-21).

**1- Mutually protected articulation (MPA)** is characterized by mutual support: posterior teeth support the anterior teeth in maximal intercuspal position (MIP) while the anterior teeth maintain a clearance of 20-30 $\mu$ m. In turn, the anterior teeth separate the posterior teeth during all excursive movements (22,23). In classic form, incisors play the role of protrusive guidance, while canines guide the laterotrusive movements (24).

**Canine protected articulation (CPA)** is one of the deviations where canine teeth separate other teeth in all excursive movements (22-24).

**Anterior protected articulation (APA)** or anterior group function is another variation where canine and another anterior tooth (mostly lateral incisor) separate the posterior teeth in all eccentric movements (24).

**2- Group function occlusion (GF)** is another occlusal scheme where a group of posterior teeth play the role of guidance in lateral movements (22-24). When the anterior teeth are periodontally compromised, or could not guide the occlusion (e.g. anterior open bite or cross bite), the anterior-most posterior tooth plays the role of protrusive guidance (24).

**3- Bilateral balanced articulation (BBA)** is the preferred scheme for stabilizing removable full dentures in one or both arches. In maximal intercuspal position, there is no contact on anterior teeth. In eccentric movements, the teeth (two teeth or more in each side) share bilateral contacts in laterotrusive, or antero-posterior contacts in protrusive movements (22,23).

**Lingualized articulation** is one of subgroups where lingual cusps of maxillary teeth play the main role of centric and eccentric contacts to improve the load distribution and direction (25).

Implant-specified occlusion or implant protected occlusion (IPO), introduced by Misch (1) considers dental implant requirements in designing occlusal schemes. The differences mainly return to the lack of periodontal ligament in implants (1-6,22) that necessitates decreasing the occlusal loads and directing the forces along implants' long axes (1-6). The principles of implant protected occlusion could be summarized in the following considerations: 1- Any occlusal plane inconsistency should be corrected before impression (1).

2- Contacts should be centered in maximal intercuspation, with 1-1.5 mm freedom in centric (1,5,19,22).

3- Occlusal timing: In maximal intercuspal position there should be no or light contacts on

implants during light tapping, and identical contact on implant and adjacent teeth in heavy tapping (1,3,5,6,8,26). This principle should also be followed in eccentric movements where adjacent teeth are the main guidance and implant-contact happens just after completion of lateral movement of natural teeth. Timing returns to the difference between natural movability of implant and tooth.

4- The occlusal table should be narrower for implant restorations by reducing palatal contour of maxillary and buccal contour of mandibular restorations (1,5,8,15,22,26,27).

5- Incisal guidance angulation should be reduced to 23- 25 degrees on implants (5,21-23,28).

6- The cuspal inclination of posterior implant restorations should be reduced (1,5,8,15).

7-Point contacts particularly cusp-fossa occlusion is preferred on implant restorations (5,15).

After implant treatment, the patient's previous occlusion might be maintained or changed based on the number of replaced teeth, oral conditions, and occlusal stability (29-31). Considering the implants' characteristics and based on scientific evidence, it is preferred to disclude (D) implants by natural teeth contacts in all excursive movements, however, if this ideal scenario is not applicable, sharing (S) the discluding contacts on teeth and implants (with attention to timing in lateral movements), or splinted (S) implants could provide more predictable long-term results (1-3,5,6,7,14,15, 32). This concept will be referred to as DSS rule to facilitate explanation. DSS rule stands for: disclude the implants by natural teeth share the discluding contacts between natural teeth and implants, or splint the implants to predictably sustain the discluding role. Fixed implant restorations will be classified into single tooth, fixed partial restorations, full arch, and full mouth reconstructions to facilitate describing the preferred occlusion for each situation.

#### **A. Single tooth implant restorations**


For all single implant restorations, occlusal timing is necessary in maximal intercuspal position. In eccentric movements, the loads should be mainly sustained by adjacent teeth and patient's existing occlusion should be followed (1).

**A-1. Posterior or anterior single-tooth implant:** Implant restoration should preferably be discluded during all eccentric movements, or participate in eccentric guidance, if necessary, following timing principle in lateral movements (3,5,6,8,26,27). The intensity of occlusal contacts could be gradually increased within 3 months intervals after restoration insertion (33), since the occlusal force and contact time of implant prostheses will change significantly with time (34).

**A-2. Canine single-tooth implant:** Canine single implant is preferred not to burden the whole guide role in canine-raised occlusion (1). The preferred occlusion is mutually protected articulation (anterior protected) or group function, without canine participation (2,6,26). An experimental study has shown that group function occlusion produces less strain in canine implant restorations (35). In cases of inevitable canine engagement, the dentist should attempt to include at least one natural tooth, and take the advantage of natural proprioception (1,36). Anterior teeth, e.g. lateral incisors, are preferred since they are farther from the temporomandibular joint (1,24). During light lateral excursion, the periodontally healthy lateral incisor occludes first and moves 97  $\mu$ m. Afterwards, canine implant engages and participates in posterior teeth disclusion (1). During heavy excursion, the natural teeth and implant crown contact with similar magnitude (1). However, in Angle's skeletal class II-division 1, anterior crossbite or open bite, the first premolar may need to be participated instead of lateral incisor (1,26).

#### **B. Partially edentulous patients**

For all fixed implant restorations inserted in partially edentulous jaw, timing is required in maximal intercuspal position or centric relation (1,32). In all eccentric movements, disclusion of implants is preferred. However, in inevitable participation, implants should follow the guidance provided by adjacent teeth under the timing principles, or be splinted together (DSS rule) (1,5,8,32). Although first proposed for removable denture, Kennedy classification will be used in the following section to discriminate different situations (See figure 2).



**Figure 2.** The examples of some different types of Kennedy classification: A. Cl I extended to anterior with anterior modification, B. Cl II extended to anterior with anterior modification, C. posterior cl III with posterior modification

**B-1. Kennedy Class I:** (bilateral posterior edentulism): When only posterior teeth are substituted by implants, mutually protected articulation on natural dentition is strongly preferred (3,6,27). All movements are preferred to be guided by anterior teeth, and there should be no interference on implants in centric relation or eccentric movements (1,5,8). However, when posterior edentulism has been extended to anterior, or situations and prognoses of remaining anterior teeth do not let them separate implant restorations independently, the participation between teeth and implants could be helpful, and anterior and posterior group function could be used for protrusive and laterotrusive movements, subsequently (5,6,8,15,22,26,27,32). In extended posterior edentulism to anterior, implants' splinting or non-splinting will determine the occlusion. When the implants are splinted

together, mutually protected articulation is the ideal occlusion. When the implants are not splinted, group function or mutually protected articulation following DSS rule could be considered (5,6,8,15,22,26,27). When only anterior teeth are remained, the amount of centric relation clearance between anterior teeth in mutually protected articulation should be less than normal value (20-30  $\mu\text{m}$ ) since posterior implants are responsible for that, however, posterior clearance provided by natural dentition could be the same as normal or less, depending on remaining teeth situations (36-39). It has also been suggested to use cross bite occlusion in posterior implant restorations to minimize off-axis loads on implants (1,5,8,22,26).


**B-2. Kennedy Class II:** The same principles as class I are applied for unilateral posterior edentulism.

**B-3. Kennedy Class III:** In this type (anterior or posterior edentulous areas surrounded by teeth), DSS rule is followed, with special emphasis on the priority of implant disclusion (32). The disclusion amount is determined by natural occlusal characteristics (6,27). However, 1.1mm posterior disclusion in protrusion, and 0.5 (for working side) to 1mm (for non-working side) clearance in lateral movements have been proposed as the least acceptable disclusion quantities (40,41). It is preferred to use the patient's previous occlusion. However, if canine has also been replaced by implant, the splinting condition plays the same role as Kennedy class I and II (1-9,14-24). A finite element analysis showed that using canine protected articulation in an implant bridge replacing canine and premolar teeth causes less overall stress (42).

**B-4. Kennedy Class IV:** Anterior edentulous area is restored by implant-restorations in this class. The anterior implants should have 30 $\mu\text{m}$  clearance in centric relation/maximal intercuspation, and should be supported by posterior teeth (mutually protected articulation) (6,8,26,27). The canine situation determines occlusal scheme in eccentric movements (26,43). If the canine is a natural tooth, mutually protected articulation (canine guided) could be the preferred occlusion (26,43). However, when the canine has been replaced by implant, anterior or posterior group function is preferred (2,6,22,26,43). Canine guided articulation could also be used if canine implant is splinted to the adjacent implants (26,43). Generally, the anterior guidance is

preferred over posterior guidance in all excursions (2,3,8,26), provided that anterior implants are not overloaded (26,43). For this purpose, anterior guidance must be flatter than normal (less vertical, and more horizontal anterior overlap) (3,8,26,27), the amount of freedom in centric occlusion must be greater than normal (3), and the amount of disclusion should be less (0.8 mm for protrusive movement, and 0 (working side) to 0.4 mm (non-working side) for lateral excursions) (2).

### Occlusal schemes for modification parts of partially edentulous patients (Figure 3)



**Figure 3.** Occlusal schemes for modification parts in partially edentulous patients. MPA: mutually protected articulation, GF: group function occlusion, APA: anterior protected articulation, CPA: canine protected articulation

The edentulous areas separated from the main part will be referred to as “modifications” based on Kennedy’s classification. Modifications are located more anterior than the main section, and therefore, they could be present in Kennedy’s class I, II, or III, but not in class IV (44). The implant-supported fixed prostheses in modifications might have their own requirements and considerations for occlusion and there are situations where the occlusions have to be different in two sides of the arch. If a modification section is located in posterior part of the mouth, the ideal occlusal scheme will be mutually protected articulation on natural anterior teeth (2,5-8). However, for anterior modifications, canine situation is determinative. If the canine is not replaced by implant,

mutually protected articulation (canine guided or classic type) is preferred (9,14,15). When canine is replaced by implant and the implants are splinted, mutually protected articulation (all types) could be an ideal occlusion. When the implants are not splinted in anterior canine containing modification part, group function occlusion and mutually protected articulation (anterior protected type) both have been suggested in different studies (20-23). In recent researches, mutually protected articulation is preferred not only for fixed restorations, but also for removeable dentures with evidences of improved masticatory efficacy, patient satisfaction, and reduced masticatory muscles activity that decrease the occlusal loading on the whole dentition (Table 1) (45-49). Therefore, this occlusal pattern will be preferred wherever it could be applied, provided that single implant does not guide the eccentric movements separately and DSS rule is followed for more predictability (2,3,5-9,14-16). There are limited evidence-based documents on occlusal scheme for modification parts of partially edentulism. These limitations have been considered in suggesting occlusal schemes in table 1, and strongly call for further studies.

### C. Fixed full arch implant restorations

Timing in centric relation is not applicable when the whole arch has been restored by implants (1). The ideal occlusal scheme will be determined according to the opposing arch situations. When the implant prosthesis is opposed by removable full denture, bilateral balanced lingualized occlusion is preferred (5,8,15,22,26,50,51). However, recent studies have clarified that mutually protected articulation “promotes self-perception and has positive impact on masticatory function” even in complete denture wearers, that requires further evaluative researches (46,47). In the presence of removable denture in the opposing arch, maximal intercuspal position and centric relation should be coincident with no interferences in functional movements (5). To reduce the effect of non-axial forces, 1-1.5 mm freedom in centric has been suggested (5,22), the anterior guidance should be shallow (22), and identical intensity contacts in centric

**Table 1.** Summary of articles on comparing canine protected articulation (CPA) and bilateral balanced articulation (BBA) in removable prostheses.

Reference	Prosthesis type	Comparison	Duration	Conclusion
El-sadany HF, 2020 <sup>(45)</sup>	Complete denture	BBA vs. CPA	1 months	Both occlusal schemes could be used effectively with no significant difference in masticatory function.
Brandt S, 2019 <sup>(46)</sup>	Complete denture	BBA vs. CPA	3 months	CPA occlusion offered significantly better retention, mastication, phonetics and esthetics.
Pero AC, 2019 <sup>(47)</sup>	Complete denture	BBA vs. CPA	1 month	CPA promoted self-perception of the patient, their ability to chew certain foods, and masticatory function
Lemos CAA, 2018 <sup>(48)</sup>	Complete denture	Different occlusal schemes	Systematic review	CPA reduces muscular activity and should be used for complete denture.
Greco GD, 2008 <sup>(49)</sup>	Implant supported mandibular overdenture (RP5)	Different occlusal schemes	In-vitro	CPA should be the pattern of choice. BBA should not be applied.

relation and all movements should be considered on maximum participating implants (6,8,26). In the presence of natural dentition or fixed implant restorations in opposing arch, the ideal occlusion is mutually protected articulation. However, some researchers proposed group function occlusion (5,8,15,22, 26,52,53). The possible reason might be the angulation and smaller diameter of anterior implants, which make them more susceptible to loading during excursive movements. Nevertheless, according to a finite element analysis study, even in such situations, canine protected articulation produces less stress than group function (54).

#### **D. Fixed full mouth implant rehabilitation**

Timing in centric is not applicable, and mutually protected articulation is the option of choice for eccentric guidance in full fixed implant rehabilitations (2,6). Coincident centric relation and maximal intercuspal position (6), 30µm clearance for anterior restorations in centric relation, and shallow anterior guidance (2) should be considered. The suggested amount of disclusion is 1mm for protrusive movements,

and 0.3 mm (in working side) to 0.8 mm (in non-working side) for laterotrusive movements (2).

There is conflicting evidence regarding the ideal occlusal scheme for full mouth implant restorations (55). According to the Miralles, canine-protected articulation has the same stress profile as group function, and both are acceptable schemes. However, a finite element analysis showed that group function is not appropriate for high stress production, while canine protected articulation causes the lowest stress value during lateral movements (56). It is a reasonable occlusal scheme for full mouth fixed implant restorations.

Occlusion is an important determinative factor for long-lasting success of implant treatments. Selection of appropriate occlusal scheme in each of the implant prosthetic options is challenging. This review study tried to gather available diverse evidences to facilitate this decision-making challenge. Table 2 shows the summary of occlusal considerations in implant fixed prosthetics.

**Table 2.** Summary of articles on occlusal considerations in fixed implant restorations. BBA: bilateral balanced articulation, MPA: mutually protected articulation, CPA: canine protected articulation, GF: group function occlusion.

Reference	Prosthesis type	Comparison	Study type	Conclusion
<b>Seifi M, 2017</b> (61)	Full mouth fixed implant restorations	GF vs. MPA	In vivo: Not mentioned	MPA showed lower Masseter and Temporalis muscles contraction
<b>Hasan I, 2014</b> (59)	Long span cement able fixed implant prosthesis	Splinting vs. non-splinting	In-vitro: finite element analysis	Splinting reduced the stresses on cortical bone
<b>Gore E, 2014</b> (64)	Fixed partial denture supported by implant	GF vs. CPA	In-vitro: finite element analysis	CPA had less functional loading on implants
<b>Teixeira FM, 2012</b> (58)	2 implants in second premolar and molar	Different force magnitude: 300,600 and 900 N	In-vitro: photoelastic model	Splinting the crowns of adjacent implants decreases the stress especially in 600 N of load.
<b>Yokoyama S, 2005</b> (62)	Mandibular bone supporting a single or separate multiple implant-superstructures	Splinting vs. non-splinting CPA vs. GF	In-vitro: finite element analysis	Splinted crowns had lower stresses in the peri-implant bone. CPA had lower stresses in working side implants.
<b>Guichet DL, 2000</b> (57)	Implant supported crowns	Splinting vs. non-splinting	In-vitro: photoelastic model	Splinted restorations exhibited better load sharing compared to non-splinted ones.
<b>Apicella A, 1998</b> (60)	Mandibular fixed full arch	MPA vs. GF	In-vitro: finite element analysis	GF had less stress intensity on the cortical bone
<b>Hobkirk JA, 1996</b> (63)	Mandibular implant supported bridges (All-on-3, All-on-4)	GF vs. BBA	In vivo: 6 months	BBA had lower mean peak masticatory force

All the occlusion terms have been derived from the latest version of the Glossary of Prosthodontic Terms to facilitate understanding (65).

Although preferred occlusal scheme for each prosthetic situation has been proposed based on available literature and evidence, there is a severe lack in clinical studies or even evaluative in-vitro researches on long-term effects of these occlusal schemes on implant, restorations, and surrounding tissues. Since occlusal loading plays an effective role in implant successful serviceability, evaluative studies on this basic

other encountered limitations were the inconsistency and incoherence in review articles, inadequate studies on this topic, lack of clear guidelines for choosing appropriate implant occlusion in different clinical situations, the complexity of occlusion concept, and the differences between implant and tooth occlusal requirements. The present review tried to collect available scientific documents and help the clinicians organize the diverse data for sound decision making on occlusal scheme selection. However, this review focused on normal situations, while there are several

compromised situations that call for further studies. Abnormal occlusal patterns namely excessive horizontal or vertical overlap, Angle's class II or III classification, cross-bite, or open-bite may affect the selection criteria of occlusion as well as abnormal conditions such as periodontally compromised teeth, horizontal or vertical cantilever on implants, and patient special parafunctional habits. All these fields could be attractive goals for researches warranting further studies.

### Conclusion

Occlusion plays an important role in prosthetic and implant complications and failures. Considering the differences between implant and natural tooth, occlusion could even be more determinative in long-term serviceability of implant-based fixed prostheses. DSS rule (disclusion, sharing, splinting) has been introduced based on available evidence to facilitate decision making in a variety of clinical situations. Understanding and applying correct occlusal principles and patterns are critically important for any clinician, especially given the increasing use of implant restorations.

### References

- Misch CE. Dental implant prosthetics. 2<sup>nd</sup> edition. Missouri, Elsevier Mosby 2015; ch1: pp1-25, ch24: pp 600-614, ch30: pp 829-873, ch31: pp 874-912.
- Jacob S, Nandini V, Nayar S, Gopalakrishnan A. Occlusal principles and considerations for the osseointegrated Prosthesis. J of Dental and Medical Sciences 2013;3(5):47-54.
- Davies SJ, Gray RJ, Young MP. Good occlusal practice in the provision of implant borne prostheses. British Dental Journal 2002; 192 (2):79-88.
- Simonis P, Dufour T, Tenebaum H. Long-term implant survival and success: a10-16-years follow-up of non-submerged dental implants. Clin Oral Impl Res 2010;21(7):772-777.
- Kim Y, Oh TJ, Misch CE, Wang HL. Occlusal considerations in implant therapy: clinical guidelines with biomechanical rationale. Clin Oral Impl Res 2005;16(1):26-35.
- Rilo B, da Silva JL, Mora MJ, Santana U, Compostela S. Guidelines for occlusion strategy in implant-borne prostheses- A review. International Dental Journal 2008;58(3):139-145.
- Naert I, Quirynen M, van Steenberghe D, Darius P. A study of 589 consecutive implants supporting complete fixed prostheses. Part II: Prosthetic aspects. J Prosthet Dent 1992;68(6): 949-56.
- Abichandani SJ, Bhojaraju N, Guttal S, Srilashmi J. Implant protected occlusion: A comprehensive review. European Journal of prosthodontics 2013; 1 (2):29-36
- Schwarz MS. Mechanical complications of dental implants. Clin Oral Implants Res 2002; 11(s1):156-158.
- Sadowsky SJ. Occlusal overload with dental implants: a review. Int J Implant Dent 2019; 5 (1):29.
- Isidor F. Loss of osseointegration caused by occlusal load of oral implants. A clinical and radiographic study in monkeys. Clin Oral Implants Res 1996;7(2):143-52.
- Isidor F. Histological evaluation of peri-implant bone at implants subjected to occlusal overload or plaque accumulation. Clin Oral Implants Res 1997; 8 (1):1-9.
- Isidor F. Clinical probing and radiographic assessment in relation to the histologic bone level at oral implants in monkeys. Clin Oral Implants Res 1997;8(4):255-64.
- Verma M, Nanda A, Sood A. Principles of occlusion in implant dentistry. Journal of the international clinical dental research organization 2015;7(1):27-33.
- Chen YY, Kuan CL, Wang YB. Implant occlusion: biomechanical considerations for implant-supported prostheses. J Dent Sci 2008; 3(2):65 -74.
- Gartner JL, Mushimoto K, Weber HP, Nishimura I. Effect of osseointegrated implants on the coordination of masticatory muscles: a pilot study. J Prosthet Dent 2000; 82(4):185-193.
- Adell R, Lekholm U, Rockler B, Branemark PI. A 15-year study of osseointegrated implants in the treatment of the edentulous jaw. Int J Oral Surg 1981;10(6):387-416.
- Curtas S, Chapman G, Meguid MM. Evaluation of nutritional status. Nurs Clin North Am 1989; 24(2): 301-313.

19. Lundgren D, Laurell L. Biomechanical aspects of fixed bridgework supported by natural teeth and endosseous implants. *Periodontol* 2000; 4:23-40.
20. Wismeijer D, van Waas MA, Kalk W. Factors to consider in selecting an occlusal concept for patients with implants in the edentulous mandible. *J Prosthet Dent* 1995;74(4):380-384.
21. Mericske-Stern RD, Taylor TD, Belser U. Management of the edentulous patient. *Clin Oral Implants Res* 2000;11(s1):108-125.
22. Jambhekar S, Kheur M, Kothavade M, Dugal R. Occlusion and Occlusal Considerations in Implantology. *Indian journal of dental advancements* 2010;2(1):125-130.
23. Shilinburg HT, Sather DA, Wilson EL, Cain JR, Mitchell DL, Blanco LJ, et.al. *Fundamentals of fixed prosthodontics*. 4<sup>th</sup> ed. Quintessence Publishing, Hanover; 2013: ch 2: pp 13-25.
24. Dawson PE. *Functional occlusion from TMJ to smile design*. Philadelphia: Linda Duncan; 2007, ch 21: pp 207-230.
25. Becker CM, Swoope CC, Guckes AD. Lingualized occlusion for removable prosthodontics. *J Prosthet Dent* 1977; 38(6): 601-608.
26. Gross MD. Occlusion in implant dentistry: A review of the literature of prosthetic determinants and current concepts. *Australian Dental Journal* 2008; 53(1):60-68.
27. Yuan JC, Sukotjo C. Occlusion for implant-supported fixed dental prostheses in partially edentulous patients: a literature review and current concepts. *J Periodontal Implant Sci* 2013; 43 (2):51-57.
28. Wismeijer D, Van Waas MA, Kalk W. Factors to consider in selecting an occlusal concept for patients with implants in the edentulous mandible. *J Prosthet Dent* 1995;74(4):380-384.
29. Foster LV. Clinical aspects of occlusion: Occlusal terminology and the conformational approach. *Dent Update* 1992;19(8):345-348.
30. Ash MM, Ramfjord SP. *Occlusion*. 2<sup>nd</sup> ed. Philadelphia: Saunders; 1971. pp 1178-9.
31. Davies S, Gray RMJ. What is occlusion? *Br Dent J* 2001;191(5):235-245.
32. Hobkirk JA. *Functional occlusion in restorative dentistry and prosthodontics*. Elsevier Mosby 2016; ch16: pp143-151.
33. Madani AS, Nakhaei M, Moazzami SM, Alami M, Rajati Haghi H, Moazzami SM. Post-insertion posterior single-implant occlusion changes at different intervals: A T-Scan computerized occlusal analysis. *J Contemp Dent Pract* 2017; 18 (10):927-932.
34. Luo Q, Ding Q, Zhang L, Xie Q. analyzing the occlusion variation of single posterior implant-supported fixed prostheses by using the T- scan system: a prospective 3-year follow up study. *J Prosthet Dent* 2020;123(1):79-84.
35. Lu J, Abduo J, Palamara J. Effect of different lateral occlusion schemes on peri-implant strain: a laboratory study. *J Adv Prosthodont* 2017;9(1):45-51.
36. Ishigaki S, Nakano T, Yamada S, Nakamura T, Takeshima F. Biomechanical stress in bone surrounding an implant under simulated chewing. *Clin Oral Impl Res* 2003;14(1):97-102.
37. Engquist B, Bergendal T, Kallus T. A retrospective multicenter evaluation of osseointegrated implant supporting overdentures. *Int J Oral Maxillofac Implants* 1988;3(2):129-134.
38. Jemt T, Chai J, Harnett J. A 5- year prospective multicenter follow-up report on overdentures supported by osseointegrated implants. *Int J Oral Maxillofac Implants* 1996; 11(3):291-298.
39. Wismeijer D, Van Maas MAJ, Vermeeren J. Overdentures supported by implants: A 6.5-year evaluation of patient satisfaction and prosthesis aftercare. *Int J Oral Maxillofac Implants* 1995; 10 (6):744-749.
40. Jemt T, Stallard PA. The effect of chewing movements on changing mandibular complete dentures to osseointegrated overdentures. *J Prosthet Dent* 1986;55(3):357-361.
41. Geertman ME, Slagter AP, Van Maas MAJ. Commintion of food with mandibular implant-retained overdentures. *J Dent Res* 1994; 73(12): 1858-1864.
42. Anaraki MR, Torab A, Mounesi Rad T. Comparison of stress in implant-supported monolithic zirconia fixed partial dentures between canine guidance and group function occlusal patterns: a finite element analysis. *J Dent Res Dent Clin Dent Prospects* 2019; 13(2): 90-97.
43. GrossmannY, Finger IM, Block MS. Indications for splinting implant restoration. *J Oral Maxillofac Surg* 2005;63(11):1642-52.
44. Phoenix RD, Cagna DR, Defreest CF. *Stewart's Clinical removable partial prosthodontics*. 4<sup>th</sup> ed.

- Quintessence Publishing, Hanover; 2008, ch 1: pp 10.
45. El-sadany HF, El-fattah A, El-Mahrouky NA, Kabeel SM. The effect of different occlusal concepts on masticatory efficiency in implant supported mandibular over-dentures. *Al-Azhar Dental Journal for girls* 2020;7(1):41-48.
46. Brandt S, Danielczak R, Kunzmann A, Lauer HC, Molzberger M. Prospective clinical study of bilateral balanced occlusion versus canine-guided occlusion in complete denture wearers. *Clinical Oral Investigations* 2019; 23 (11):4181-4188.
47. Pero AC, Scavassin PM, Policastro VB, de Oliveira Junior NM, Mendoza Marin DO, Silva MDDD, et al. Masticatory function in complete denture wearers varying degree of mandibular bone resorption and occlusion concept: canine guide occlusion versus bilateral balanced occlusion in a cross-over trial. *J prosthodont res* 2019;3(5):552-559.
48. Lemos CAA, Verri FR, Gomes JML, Santiago Junior JF, Moraes SLD, Pellizzer EP. Bilateral balanced occlusion compared to other occlusal schemes in complete dentures: A systematic review. *J Oral Rehabil* 2018;45(4):344-354.
49. Greco GD, Jansen WC, Lander Junior J, Seraidarian PI. Biomechanical analysis of the stresses generated by different disocclusion patterns in an implant supported mandibular complete dentures. *J Appl Oral Sci* 2009; 17(5): 515-520.
50. Bonwill WGA. The geometric and mechanical laws of the articulation of the human teeth, The anatomical articulator. Philadelphia, Lea Brothers;1887.
51. Spee FG. The condylar path of the mandible along the skull. *Arch Anat Physiol* 1890;16(1): 285-94.
52. Stuart CE, Stallard H. Principles involved in restoring occlusion to natural teeth. *J Prosthet Dent* 1960;10(2): 304-313.
53. D'Amico A. Functional occlusion of the natural teeth in man. *J Prosthet Dent* 1961; 11 (5):899-915.
54. Yokoyama S, Wakabayashi N, Shitoa M, Ohayama T. Stress analysis in edentulous mandibular bone supporting implant-retained 1-piece or multiple superstructures. *Int J Oral Maxillofac Implants* 2005; 20(4):578-83.
55. Miralles R. Canine-guide occlusion and group function occlusion are equally acceptable when restoring the dentition. *J Evid Based Dent Pract* 2016;16(1):41-43.
56. Turker N, Buyukkaplan US, Sadowsky SJ, Ozarslan MM. finite element stress analysis of applied forces to implants and supporting tissues using the all on four concepts with different occlusal schemes. *J Prosthodont* 2019; 28(2):185-194.
57. Guichet DL, Yoshinobu D, Caputo AA. Effect of splinting and interproximal contact tightness on load transfer by implant restorations. *J Prosthet Dent* 2002;87(5):528-535.
58. Teixeira FM, de Assis Claro CA, Neves AC, de Mello Rode S, da Silva-Concilo LR. Influence of loading and use of occlusal splint in implant-supported fixed prostheses. *Journal of Craniofac Surg* 2012; 23 (5):477-480.
59. Hasan I, Bourauel C, Keilig L, Stark H, Luckerath W. The effect of implant splinting on the load distribution in bone bed around implant-supported fixed prosthesis with different framework materials: A finite element study. *Annals of Anatomy-Anatomischer Anzeiger* 2015; 199:43-51.
60. Apicella A, Masi E, Nicolais L, Zarone F, Rosa N DE, Valletta G. A finite-element model study of occlusal schemes in full-arch implant restoration. *J Mater Sci Mater Med*1998; 9(4): 191-196.
61. Seifi M, Nodehi D, Ghahramanloo A, Ahmadi Z, Farhangnia A, Saedi M, et al. Comparing the Electromyographic features of the Masseter and Temporal muscles in patients with full mouth implant-supported FDPs and natural dentition. *Journal of Academy of medical science of Bosnia and Herzegovina* 2017;25(1):49-53.
62. Yokoyama S, Wakabayashi N, Shiota M, Ohyama T. Stress analysis in edentulous mandibular bone supporting implant-retained 1-piece or multiple superstructures. *Int J Oral Maxillofac Implants* 2005;20(4):578-583.
63. Hobkirk JA, Brouziotou-Davas E. The influence of occlusal scheme on masticatory forced using implant stabilized bridges. *J of Oral Rehab* 1996; 23:386-391.
64. Gore E, Evlioglu G. Assessment of the effect of two occlusal concepts for implant-supported fixed prostheses by finite element analysis in patients with bruxism. *J Oral Implantol* 2014; 40 (1):68-75.
65. Driscoll CF, Freilich MA, Guckes AD, Knoernschild KL, McGarry TJ. Glossary of Prosthodontic Terms Committee of the Academy of Prosthodontics. *J Prosthet Dent* 2017;117(5S):e1-e10.